

Healthcare Effectiveness Data and Information Set (HEDIS[®])

2011 Measures

Adolescent Well Care

The percentage of enrolled members 12-21 years of age who had at least one comprehensive well-care visit with a PCP or OB/GYN practitioner during the measurement year.

Adults' Access to Preventive/Ambulatory Health Services

The percentage of members 20 years and older who had an ambulatory or preventive care visit.

Ambulatory Care

The summary of utilization of ambulatory care for outpatient visits and ED visits.

Annual Dental Visits

The percentage of members age 2-21 who had at least one dental visit during the measurement year.

Antidepressant Medication Management- Acute Phase Treatment

The percentage of members 18 years of age and older who were diagnosed with a new episode of major depression and remained on antidepressant medication for at least 84 days.

Antidepressant Medication Management- Continuation Phase Treatment

The percentage of members 18 years of age and older who were diagnosed with a new episode of major depression and remained on antidepressant medication for at least 180 days.

Adult BMI Assessment

The percentage of members 18-74 years of age who had an outpatient visit and their body mass index documented during the measurement year.

Breast Cancer Screening

The percentage of women 40-69 years of age who had a mammogram to screen for breast cancer during the measurement year and the year prior to the measurement year.

Cervical Cancer Screening

One or more Pap tests during the measurement year or the two years prior to the measurement year.
Women age 24 – 64

Chlamydia Screening for Women

The percentage of women 16–24 years of age who were identified as sexually active and who had at least one test for chlamydia during the measurement year.

Childhood Immunization Status

The percentage of children 2 years of age who had four diphtheria, tetanus and acellular pertussis (DTaP); three polio (IPV); one measles, mumps and rubella (MMR); three H influenza type B (HiB); three hepatitis B (HepB), one chicken pox (VZV); four pneumococcal conjugate (PCV); two hepatitis A (HepA); two or three rotavirus (RV); and two influenza (flu) vaccines by their second birthday.

Comprehensive Diabetes - HbA1C Testing

HbA1C test was performed during the measurement year.

Comprehensive Diabetes - HbA1C Poor Control (Inverse Measure)

The most recent HbA1C level is >9.0% or is missing or was not done during the measurement year.

Comprehensive Diabetes – HbA1C Good Control

The most recent HbA1c level is <8.0%.

Comprehensive Diabetes - Eye Exam

A retinal or dilated eye exam by an eye care professional was done during the measurement year OR a negative retinal exam was done by an eye professional in the year prior to the measurement year.

Comprehensive Diabetes - LDL Screening

An LDL-C test was performed during the measurement year.

Comprehensive Diabetes - LDL-C Control

The most recent LDL-C level performed during the measurement year is <100 mL.

Comprehensive Diabetes – Nephropathy

A urine microalbumin test was done during the measurement year OR there is evidence of nephropathy during the measurement year.

Controlling High Blood Pressure

The percentage of members 18-85 with hypertension whose most recent BP is <140/90.

Follow-up Care for Children Prescribed ADHD Medication

The percentage of children newly prescribed attention-deficit/hyperactivity disorder (ADHD) medication who had at least three follow-up care visits within a 10-month period, one of which was within 30 days of when the first ADHD medication was dispensed. Two rates are reported.

- *Initiation Phase.* The percentage of members 6–12 years of age as of the IPSD with an ambulatory prescription dispensed for ADHD medication, who had one follow-up visit with practitioner with prescribing authority during the 30-day Initiation Phase.
- *Continuation and Maintenance (C&M) Phase.* The percentage of members 6–12 years of age as of the IPSD with an ambulatory prescription dispensed for ADHD medication, who remained on the medication for at least 210 days and who, in addition to the visit in the Initiation Phase, had at least two follow-up visits with a practitioner within 270 days (9 months) after the Initiation Phase ended.

Adolescent Immunization Status

The percentage of adolescents 13 years of age who had one dose of meningococcal vaccine and one tetanus, diphtheria toxoids and acellular pertussis vaccine (Tdap) or one tetanus, diphtheria toxoids vaccine (Td) by their 13th birthday.

Lead Screening in Children

The percentage of children 2 years of age who had one or more capillary or venous blood tests for lead poisoning by their second birthday.

Mental Health Utilization-Inpatient, Intermediate, & Ambulatory Services

The number and percentage of members receiving the following mental health services during the measurement year: Any service, Inpatient, Intensive outpatient or partial hospitalization, Outpatient or ED

Pharyngitis

The percentage of children 2–18 years of age, who were diagnosed with pharyngitis, dispensed an antibiotic and received a group A streptococcus (strep) test for the episode.

Prenatal Care

The percentage of women who delivered a live infant that received a prenatal care visit as a member of the plan in the first trimester or within 42 days of enrollment in the plan.

Postpartum Care

The percentage of women who delivered a live infant that had a postpartum visit on or between 21 and 56 days after delivery.

Use of Appropriate Medications for People with Asthma

The percentage of members 5-50 years of age during the measurement year who were identified as having persistent asthma and who were appropriately prescribed medication during the measurement year.

Well-Child Visits in the first 15 months

The percentage of children who turned 15 months old during the measurement year who had six or more well-child visits with a PCP.

Well-Child Visits in the 3rd, 4th, 5th, & 6th Years of Life

The percentage of children 3-6 years of age who received one or more well-child visits with a PCP during the measurement year

Follow-Up after Hospitalization for Mental Illness

The percent of enrollees who were hospitalized for a mental health diagnosis and

were discharged to the community from an acute care facility and were seen on an outpatient basis by a mental health practitioner within seven days and within 30 days.

Mental Health Readmission Rate

The percentage of enrollees who were hospitalized for a mental health diagnosis and were discharged to the community from an acute care facility and were readmitted for a mental health diagnosis within 30 days.

Lipid Profile Annually

The percentage of members 18 to 85 years of age with a diagnosis of hypertension who had a lipid profile that includes total cholesterol, HDL, and triglycerides during the measurement year.

Use of Angiotensin-Converting Enzyme (ACE) Inhibitors/ Angiotensin Receptor Blockers (ARB) Therapy

The percentage of members 18 years and older during the measurement year who were diagnosed with congestive heart failure, and who have at least one prescription filled for ACE inhibitors or ARBs during the measurement year.

Frequency of Ongoing Prenatal Care - Greater than 81% of Visits

The percentage of Medicaid deliveries that received greater than 81% of expected prenatal visits.

AHCA: Florida Medicaid Quality in Managed Care- Performance Measure Indicator Descriptions

http://www.ahca.myflorida.com/MCHQ/Managed_Health_Care/MHMO/med_prov_0912.shtml

[National Committee for Quality Assurance \(NCQA\)](#)
